arXiv:hep-ph/0010154v2 23 Oct 2000

Functional Integral Method in the Theory of Color Superconductivity[1]

Nguyen Van Hieu
Institute of Physics, NCST of Vietnam
and
Faculty of Technology, Vietnam National University, Hanoi

Abstract

We propose to study the superconducting pairing of quarks with the formation of the diquarks as well as the quark-antiquark pairing in QCD by means of the functional integral technique. The dynamical equations for the superconducting order parameters are the nonlinear integral equations for the composite quantum fields describing the quark-quark or quark-antiquark systems. These composite fields are the bi-local fields if the pairing is generated by the gluon exchange while for the instanton induced pairing interactions they are the local ones. The expressions of the free energy densities are derived. The binding of three quarks is also discussed.

The superconducting pairing of quarks due to the gluon exchange in QCD with the formation of the diquark condensate was proposed by Barrois[1] and Frautshi[2] since more than two decades and then studied by Bailin and Love[3], Donoglue and Sateesh[4], Iwasaki and Iwado[5]. Recently, in a series of papers by Alford, Rajagopal and Wilczek[6], Schäfer and Wilczek[7], Rapp, Schäfer, Shuryak and Velkovsky[8], Evans, Hsu and Schwetz[9], Son[10], Carter and Diakonov[11] and others[12−21] there arose a new interest to the existence of the diquark Bose condensate in the QCD dense matter-the color superconductivity. The connection between the color superconductivity and the chiral phase transition in QCD was studied by Berges and Rajagopal[22], Harada and Shibata[23]. There exists also the spontaneous parity violation, as it was shown by Pisarski and Rischke[24].

For the study of many-body systems of relativistic particles with the internal degrees of freedom as well as with the virtual creation and annihilation of the particle-antiparticle pairs the functional integral technique is a powerful mathematical tool. This method was applied to the study of the color superconductivity as well as the quark-antiquark pairing in a series of papers of the group of authors in Hanoi[25−28]. In my talk I present a review of these works.

Denote ψ_A the quark field, where $A = (\alpha, a, i)$ is the set consisting of the Dirac spinor index $\alpha = 1, 2, 3, 4$ and the color and flavor indices $a = 1, 2, 3...N_c$ and $i = 1, 2, 3, ...N_f$. The internal symmetry groups are assumed to be $SU(N_c)_c$ and $SU(N_f)_f$.We work in the imaginary time formalism and denote $x = (\tau, \mathbf{x})$ a vector in the Euclidean four-dimensional space-time. The partition function of the system of free quarks and antiquarks with the chemical potential μ and at the temperature T can be expressed in the form of the functional integral

$$Z_0 = \int [D\psi] \left[D\overline{\psi}\right] \exp\left\{-\int dx \overline{\psi}^A(x)\, D_A^B \psi_B(x)\right\} \tag{1}$$

---

[1] Invited talk at the International Conference on Physics at Extreme Energy-Rencontres du Vietnam IV, Hanoi 19-25 July 2000

with brief notations

$$\int dx = \int\limits_0^\beta d\tau \int d\mathbf{x}, \qquad \beta = \frac{1}{kT}, \qquad D_A^B = \left[\gamma_4\left(\frac{\partial}{\partial\tau} - \mu\right) + \boldsymbol{\gamma}\boldsymbol{\nabla} + M\right]_A^B, \tag{2}$$

where k is the Boltzmann constant and M is the bare quark mass. We start from the study of the system with the direct instanton induced four-fermion couplings and then consider the case of the pairing due to the gluon exchange.

We write the expression of the interaction Lagrangian of the direct four-fermion couplings in two different forms

$$L_{int} = \frac{1}{2}\overline{\psi}^A(x)\overline{\psi}^C(x)V_{CA}^{BD}\psi_D(x)\psi_B(x) \tag{3}$$

and

$$L_{int} = \frac{1}{2}\overline{\psi}^A(x)\psi_B(x)U_{AC}^{BD}\overline{\psi}^C(x)\psi_D(x) \tag{4}$$

convenient for the study of the quark-quark or quark-antiquark pairing, resp. The notations V_{CA}^{BD} and U_{AC}^{BD} are related each to other

$$V_{CA}^{BD} = U_{AC}^{BD}.$$

These coupling constants are antisymmetric under the interchanges of the indices.

In order to study the quark-quark pairing we use the interaction Lagrangian in the form (3) and have the partition function

$$\begin{aligned} Z &= \int [D\psi][D\overline{\psi}] \exp\left\{-\int dx \overline{\psi}^A(x) D_A^B \psi_B(x)\right\} \\ &\quad .\exp\left\{\frac{1}{2}\int dx \overline{\psi}^A(x)\overline{\psi}^C(x)V_{CA}^{BD}\psi_D(x)\psi_B(x)\right\}. \end{aligned} \tag{5}$$

Introduce the composite local bi-spinor fields $\Phi_{DB}(x), \overline{\Phi}^{AC}(x)$ describing the diquarks and consider the functional integral

$$Z_0^{\Phi,\overline{\Phi}} = \int [D\Phi][D\overline{\Phi}] \exp\left\{-\frac{1}{2}\int dx \overline{\Phi}^{AC}(x) V_{CA}^{BD} \Phi_{DB}(x)\right\}. \tag{6}$$

Applying the corresponding Hubbard-Stratonovich transformation, we rewrite Z in the form of the functional integral over these composite local fields

$$Z = \frac{Z_0}{Z_0^{\Phi,\overline{\Phi}}} \int [D\Phi][D\overline{\Phi}] \exp\left\{S_{\text{eff}}[\Phi,\overline{\Phi}]\right\}. \tag{7}$$

The field equations

$$\frac{\delta S_{\text{eff}}[\Phi,\overline{\Phi}]}{\delta \overline{\Phi}^{AC}(x)} = 0 \tag{8}$$

are the dynamical equations for the quark-quark pairing. They can be written in the following manner

$$\begin{aligned}
\Delta_{D_1B_1}(x_1) \quad = \quad & V^{A_1C_1}_{D_1B_1}\Big\{\int dx_2 S^{C_2}_{C_1}(x_1-x_2)\Delta_{C_2A_2}(x_2)S^{A_2}_{A_1}(x_1-x_2) \\
& -\int dx_2\int dx_3\int dx_4 S^{C_2}_{C_1}(x_1-x_2)\Delta_{C_2A_2}(x_2)S^{A_2}_{A_3}(x_3-x_2) \\
& .\,\overline{\Delta}^{A_3C_3}(x_3)\,S^{C_4}_{C_3}(x_3-x_4)\,\Delta_{C_4A_4}(x_4)\,S^{A_4}_{A_1}(x_1-x_4)+....,\Big\}
\end{aligned} \tag{9}$$

where $S^B_A(x-y)$ are the propagators of the free quarks and

$$\Delta_{CA}(x) = V^{BD}_{CA}\Phi_{DB}(x), \qquad \overline{\Delta}^{BD}(x) = V^{BD}_{CA}\overline{\Phi}^{AC}(x). \tag{10}$$

Consider the special class of the constant solutions

$$\Delta_{CA} = const, \qquad \overline{\Delta}^{BD} = const.$$

Denote $\widetilde{S}^B_A(\mathbf{p},\varepsilon_m)$ the Fourier transforms of the free quark propagators, $\varepsilon_m = \frac{\pi}{\beta}(2m+1)$, and introduce $\widehat{\Delta}$, $\widehat{\overline{\Delta}}$ and $\widehat{S}(\mathbf{p},\varepsilon_m)$ the matrices with the elements $\Delta_{CA}, \overline{\Delta}^{BD}$ and $\widetilde{S}^B_A(\mathbf{p},\varepsilon_m)$. Then we have following matrix equation for the quark-quark pairing

$$\begin{aligned}
\left[\widehat{\Delta}\right]_{DB} \quad = \quad & V^{AC}_{DB}\frac{1}{\beta}\sum_m \frac{1}{(2\pi)^3}\int d\mathbf{p} \\
& \left[\frac{1}{1+\widehat{S}(\mathbf{p},\varepsilon_m)\widehat{\Delta}\widehat{S}^T(-\mathbf{p},-\varepsilon_m)\widehat{\overline{\Delta}}}S(\mathbf{p},\varepsilon_m)\widehat{\Delta}\widehat{S}^T(-\mathbf{p},-\varepsilon_m)\right]_{CA}
\end{aligned} \tag{11}$$

The corresponding free energy density equals

$$\begin{aligned}
F \quad = \quad & \frac{1}{\beta}\sum_m \frac{1}{(2\pi)^3}\int d\mathbf{p}\frac{1}{2}Tr\Big[\widehat{S}(\mathbf{p},\varepsilon_m)\widehat{\Delta}\widehat{S}^T(-\mathbf{p},-\varepsilon_m)\widehat{\overline{\Delta}}. \\
& .\left\{\frac{1}{1+\widehat{S}(\mathbf{p},\varepsilon_m)\widehat{\Delta}\widehat{S}^T(-\mathbf{p},-\varepsilon_m)\widehat{\overline{\Delta}}} - \int\limits_0^1 d\alpha \frac{1}{1+\alpha\widehat{S}(\mathbf{p},\varepsilon_m)\widehat{\Delta}\widehat{S}^T(-\mathbf{p},-\varepsilon_m)\widehat{\overline{\Delta}}}\right\}\Big].
\end{aligned} \tag{12}$$

In order to study the quark-antiquark pairing we use the interaction Lagrangian in the form (4) and start from the partition function

$$\begin{aligned}
Z \quad = \quad & \int [D\psi][D\overline{\psi}]\exp\left\{-\int dx\overline{\psi}^A(x)D^B_A\psi_B(x)\right\} \\
& .\exp\left\{\frac{1}{2}\int dx\overline{\psi}^A(x)\psi_B(x)U^{BD}_{AC}\overline{\psi}^C(x)\psi_D(x)\right\}.
\end{aligned} \tag{13}$$

Introducing the hermitian composite fields $\Phi_B^A(x)$ as well as the functional integral

$$Z_0^\Phi = \int [D\Phi] \exp\left\{-\frac{1}{2}\int dx \Phi_B^A(x) U_{AC}^{BD} \Phi_D^C(x)\right\} \tag{14}$$

and applying the corresponding Hubbard-Stratonovich transformation, we rewrite Z in the form of the functional integral over these new composite fields

$$Z = \frac{Z_0}{Z_0^\Phi} \int [D\Phi] \exp\left\{S_{\text{eff}}[\Phi]\right\}. \tag{15}$$

The field equations

$$\frac{\delta S_{\text{eff}}[\Phi]}{\delta \Phi_B^A(x)} = 0 \tag{16}$$

are the dynamical equations for the quark-antiquark pairing. They have the explicit form

$$\Delta_C^D(x) = U_{CA}^{DB} G_B^A(x,x) \tag{17}$$

where

$$\Delta_C^D(x) = U_{CA}^{DB} \Phi_B^A(x) \tag{18}$$

and $G_A^B(x,y)$ are determined by the Schwinger-Dyson equation

$$G_A^B(x,y) = S_A^B(x-y) - \int dz S_A^C(x-z) \Delta_C^D(z) G_D^B(z,y). \tag{19}$$

It is easy to verify that $G_A^B(x,y)$ are the two-point Green functions of the quarks in the presence of the quark-quark pairing. In the case of the constant order parameters

$$\Delta_A^B = const$$

the Green functions $G_A^B(x,y)$ depend only on the coordinate difference $x-y$. Their Fourier transforms $\widetilde{G}_A^B(\mathbf{p},\varepsilon_m)$ are determined by the algebraic equation

$$\widetilde{G}_A^B(\mathbf{p},\varepsilon_m) = \widetilde{S}_A^B(\mathbf{p},\varepsilon_m) - \widetilde{S}_A^C(\mathbf{p},\varepsilon_m) \Delta_C^D \widetilde{G}_D^B(\mathbf{p},\varepsilon_m). \tag{20}$$

Introducing the matrices $\widehat{G}(\mathbf{p},\varepsilon_m)$ and $\widehat{\Delta}$ with the elements $\widetilde{G}_A^B(\mathbf{p},\varepsilon_m)$ and Δ_C^D, we obtain the matrix equation

$$\frac{1}{\widehat{G}(\mathbf{p},\varepsilon_m)} = \frac{1}{\widehat{S}(\mathbf{p},\varepsilon_m)} + \widehat{\Delta}. \tag{21}$$

The free energy density of the system equals

$$\begin{aligned} F[\Delta] &= \frac{1}{\beta}\sum_m \frac{1}{(2\pi)^3}\int d\mathbf{p} Tr\Big[\widehat{\Delta}\,\widehat{S}(\mathbf{p},\varepsilon_m) \\ &\cdot\left\{\frac{1}{2}\frac{1}{1+\widehat{\Delta}(\mathbf{p},\varepsilon_m)\widehat{S}(\mathbf{p},\varepsilon_m)} - \int_0^1 d\alpha \frac{1}{1+\alpha\widehat{\Delta}(\mathbf{p},\varepsilon_m)\widehat{S}(\mathbf{p},\varepsilon_m)}\right\}\Big]. \end{aligned} \tag{22}$$

For the study of the quark-quark pairing due to the gluon exchange we start from the partition function in the form

$$
\begin{aligned}
Z &= \int [D\psi][D\overline{\psi}] \exp\left\{ -\int dx \overline{\psi}^{A}(x) D_{A}^{B} \psi_{B}(x) \right\} \\
&\quad . \exp\left\{ \frac{1}{2} \int dx \int dy \overline{\psi}^{A}(x) \overline{\psi}^{C}(y) V_{CA}^{BD}\left(x-y\right) \psi_{D}(y) \psi_{B}(x) \right\}, \qquad (23)
\end{aligned}
$$

where

$$
\begin{aligned}
V_{CA}^{BD}\left(x-y\right) &= -\frac{g^{2}}{4\pi^{2}} \sum_{I} \left(\gamma_{\mu} \otimes \lambda_{I}\right)_{A}^{B} \left(\gamma_{\mu} \otimes \lambda_{I}\right)_{C}^{D} \frac{1}{\left(x-y\right)^{2}}, \\
\left(\gamma_{\mu} \otimes \lambda_{I}\right)_{A}^{B} &= \left(\gamma_{\mu}\right)_{\alpha}^{\beta} \left(\lambda_{I}\right)_{a}^{b} \delta_{i}^{j}, \quad \sum_{I} \left(\lambda_{I}\right)_{a}^{b} \left(\lambda_{I}\right)_{c}^{d} = \frac{1}{2}\left[\delta_{c}^{b} \delta_{a}^{d} - \frac{1}{N_{c}} \delta_{a}^{b} \delta_{c}^{d} \right], \qquad (24)
\end{aligned}
$$

λ_I are the Gell-Mann matrices of the color symmetry group. In order to describe the diquark systems we introduce some bi-local composite fields $\Phi_{CD}\left(y,x\right), \overline{\Phi}^{AB}\left(x,y\right)$. Consider the functional integral

$$
Z_{0}^{\Phi,\overline{\Phi}} = \int [D\Phi][D\overline{\Phi}] \exp\left\{ -\frac{1}{2} \int dx \int dy \overline{\Phi}^{AC}(x,y) V_{CA}^{BD}\left(x-y\right) \Phi_{DB}(y,x) \right\}. \qquad (25)
$$

Applying the corresponding Hubbard-Stratonovich transformation, we rewrite the partition function in the form

$$
Z = \frac{Z_{0}}{Z_{0}^{\Phi,\overline{\Phi}}} \int [D\Phi][D\overline{\Phi}] \exp\left\{ S_{\mathrm{eff}}[\Phi,\overline{\Phi}] \right\}. \qquad (26)
$$

The field equations

$$
\frac{\delta S_{\mathrm{eff}}[\Phi,\overline{\Phi}]}{\delta \overline{\Phi}^{AC}(x,y)} = 0 \qquad (27)
$$

are the dynamical equations for the quark-quark pairing. The bi-local composite fields $\Phi_{DB}(y-x)$, $\overline{\Phi}^{AC}(x-y)$ depending only on the coordinate difference, or the linear combinations

$$
\Delta_{CA}\left(y-x\right) = V_{CA}^{BD}\left(x-y\right) \Phi_{DB}(y-x), \quad \overline{\Delta}^{BD}\left(x-y\right) = V_{CA}^{BD}\left(x-y\right) \overline{\Phi}^{AC}(x-y), \qquad (28)
$$

are the order parameters of the superconducting phase transition. Denote $\widetilde{\Delta}_{CA}\left(\mathbf{p},\varepsilon_m\right)$ and $\widetilde{\overline{\Delta}}^{BD}\left(\mathbf{p},\varepsilon_m\right)$ their Fourier transforms and $\widehat{\Delta}\left(\mathbf{p},\varepsilon_m\right)$,$\widehat{\overline{\Delta}}\left(\mathbf{p},\varepsilon_m\right)$ the matrices with these elements. Then the field equations (27) become the extended BCS equations for the color superconductivity induced by the gluon exchange

$$\begin{aligned}\widetilde{\Delta}_{CA}\left(\mathbf{p},\varepsilon_m\right) &= -g^2\sum_I\left(\gamma_\mu\otimes\lambda_I\right)_A^B\left(\gamma_\mu\otimes\lambda_I\right)_C^D.\frac{1}{\beta}\sum_n\frac{1}{\left(2\pi\right)^3}\int d\mathbf{q}\\ &\quad\frac{1}{\left(\varepsilon_n-\varepsilon_m\right)^2+\left(\mathbf{p}-\mathbf{q}\right)^2}\left[\frac{1}{1+\widehat{S}\left(\mathbf{q},\varepsilon_n\right)\widehat{\Delta}\left(\mathbf{q},\varepsilon_n\right)\widehat{S}^T\left(-\mathbf{q},-\varepsilon_n\right)\widehat{\overline{\Delta}}\left(\mathbf{q},\varepsilon_n\right)}\right.\\ &\quad\left..\widehat{S}\left(\mathbf{q},\varepsilon_n\right)\widehat{\Delta}\left(\mathbf{q},\varepsilon_n\right)\widehat{S}^T\left(-\mathbf{q},-\varepsilon_n\right)\right]_{DB}\end{aligned}\tag{29}$$

The free energy density of the system equals

$$\begin{aligned}F\left[\Delta,\overline{\Delta}\right] &= \frac{1}{\beta}\sum_m\frac{1}{\left(2\pi\right)^3}\int d\mathbf{p}Tr\left[\widehat{S}\left(\mathbf{p},\varepsilon_m\right)\widehat{\Delta}\left(\mathbf{p},\varepsilon_m\right)\widehat{S}^T\left(-\mathbf{p},-\varepsilon_m\right)\widehat{\overline{\Delta}}\left(\mathbf{p},\varepsilon_m\right)\right.\\ &\quad\left\{\frac{1}{1+\widehat{S}\left(\mathbf{p},\varepsilon_m\right)\widehat{\Delta}\left(\mathbf{p},\varepsilon_m\right)\widehat{S}^T\left(-\mathbf{p},-\varepsilon_m\right)\widehat{\overline{\Delta}}\left(\mathbf{p},\varepsilon_m\right)}-\right.\\ &\quad\left.\left.-\int_0^1 d\alpha\frac{1}{1+\alpha\widehat{S}\left(\mathbf{p},\varepsilon_m\right)\widehat{\Delta}\left(\mathbf{p},\varepsilon_m\right)\widehat{S}^T\left(-\mathbf{p},-\varepsilon_m\right)\widehat{\overline{\Delta}}\left(\mathbf{p},\varepsilon_m\right)}\right\}\right].\end{aligned}\tag{30}$$

In order to study the quark-antiquark pairing we use the partition function (23) in another form

$$\begin{aligned}Z &= \int[D\psi][D\overline{\psi}]\exp\left\{-\int dx\overline{\psi}^A(x)D_A^B\psi_B(x)\right\}\\ &\quad.\exp\left\{\frac{1}{2}\int dx\int dy\overline{\psi}^A(x)\psi_B(y)U_{AC}^{BD}\left(x-y\right)\overline{\psi}^C(y)\psi_D(x)\right\},\end{aligned}\tag{31}$$

$$U_{AC}^{BD}\left(x-y\right)=-V_{CA}^{DB}\left(x-y\right).$$

The quark-antiquark systems are described by the bi-local hermitian composite fields $\Phi_B^A\left(x,y\right)$. Introducing the functional integral

$$Z_0^\Phi=\int[D\Phi]\exp\left\{-\frac{1}{2}\int dx\int dy\Phi_B^A(x,y)U_{AC}^{BD}\left(x-y\right)\Phi_D^C(y,x)\right\}\tag{32}$$

and applying the corresponding Hubbard-Stratonovich transformation, we rewrite Z in the form

$$Z=\frac{Z_0}{Z_0^\Phi}\int[D\Phi]\exp\left\{S_{\text{eff}}[\Phi]\right\}.\tag{33}$$

The field equations

$$\frac{\delta S_{\text{eff}}[\Phi]}{\delta\Phi_D^C\left(y,x\right)}=0\tag{34}$$

are the dynamical equations for the quark-antiquark pairing

$$\Delta_C^D(y,x) = U_{CA}^{DB}(x-y)\, G_B^A(y,x)\,, \tag{35}$$

where $G_B^A(y,x)$ are determined by the Schwinger-Dyson equations

$$G_B^A(y,x) = S_A^B(y-x) - \int dx' \int dy' S_B^{A'}(y-y')\, \Delta_{A'}^{B'}(y',x')\, G_{B'}^A(x'-x) \tag{36}$$

and

$$\Delta_A^B(x,y) = U_{AC}^{BD}(x-y)\, \Phi_D^C(y,x)\,. \tag{37}$$

$G_B^A(y,x)$ are the two point Green functions of quarks in the presence of the quark-antiquark pairing.In the special class of the functions $\Delta_A^B(x-y)$ depending only on the coordinate difference these functions are the order parameters of the system. Denote $\widetilde{\Delta}_A^B(\mathbf{p},\varepsilon_m)$ their Fourier transforms and $\widehat{\Delta}(\mathbf{p},\varepsilon_m)$ the matrix with these elements. The dynamical equations (35) are

$$\begin{aligned} \widetilde{\Delta}_C^D(\mathbf{p},\varepsilon_m) \quad &= \quad g^2 \sum_I \left(\gamma_\mu \otimes \lambda_I\right)_A^D \left(\gamma_\mu \otimes \lambda_I\right)_C^B \\ &\quad \frac{1}{\beta} \sum_n \frac{1}{(2\pi)^3} \int d\mathbf{q} \frac{1}{(\varepsilon_n - \varepsilon_m)^2 + (\mathbf{p}-\mathbf{q})^2} \widetilde{G}_B^A(\mathbf{q},\varepsilon_n)\,, \end{aligned} \tag{38}$$

and the Schwinger-Dyson equations become

$$\frac{1}{\widehat{G}(\mathbf{q},\varepsilon_m)} = \frac{1}{\widehat{S}(\mathbf{q},\varepsilon_m)} + \widehat{\Delta}(\mathbf{q},\varepsilon_m)\,. \tag{39}$$

The free energy density of the system equals

$$\begin{aligned} F[\Delta] \quad &= \quad \frac{1}{\beta} \sum_m \frac{1}{(2\pi)^3} \int d\mathbf{p} Tr \Big[\widehat{\Delta}(\mathbf{p},\varepsilon_m)\, \widehat{S}(\mathbf{p},\varepsilon_m) \\ &\quad \left\{ \frac{1}{2} \frac{1}{1+\widehat{\Delta}(\mathbf{p},\varepsilon_m)\, \widehat{S}(\mathbf{p},\varepsilon_m)} - \int_0^1 d\alpha \frac{1}{1+\alpha\widehat{\Delta}(\mathbf{p},\varepsilon_m)\, \widehat{S}(\mathbf{p},\varepsilon_m)} \right\} \Big]\,. \end{aligned} \tag{40}$$

The existence of the non-vanishing order parameters Δ_{CA}, $\overline{\Delta}^{BD}$ or $\Delta_{CA}(y-x)$, $\overline{\Delta}^{BD}(x-y)$ which cannot be the singlets of the color symmetry group would mean the spontaneous breaking of the color symmetry[29].

The physical vacuum in QCD is the limiting case of the system with vanishing density ($\mu = 0$) at vanishing temperature ($T = 0$). In this limit the order parameters Δ_{DC}, $\overline{\Delta}^{AB}$ and Δ_B^A are the vacuum expectation values of the corresponding quantum composite fields - the new Higgs fields.

The functional integral method was applied also to the study of the binding of three quarks due to some effective six-fermion (non-local, in general) interactions of quarks. The systems of dynamical equations for the composite fermionic quantum triquark fields Ψ_{ABC} and $\overline{\Psi}^{ABC}$ were derived. The constant solutions of these equations in the case of the direct six-fermion couplings of quarks or their solutions depending only on the coordinate difference in the case of the non-local effective six-fermion interactions are the anticommuting order parameters.

Acknowledgements.

The author would like to express his sincere appreciation to the National Natural Sciences Council of Vietnam for the support to this work.

References.

[1]. B. C. Barrois, *Nucl. Phys.* **B129** (1977) 390.

[2]. C. Frautshi, *Asymptotic Freedom and Color Superconductivity in Dense Quark Matter, in: Proceedings of the Workshop on Hadronic Matter at Extreme Energy Density*, Ed., N. Cabibbo, Erice, Italy (1978).

[3]. D. Bailin and A. Love, *Nucl. Phys.* **B190** (1981) 175; **B190** (1981) 751; **B205** (1982) 119.

[4]. J. F. Donoghue and K. S. Sateesh, *Phys. Rev.* **D38** (1988) 360.

[5]. M. Iwasaki and T. Iwado, *Phys. Lett.* **B350** (1995) 163.

[6]. M. Alford, K. Rajagopal and F. Wilczek , *Phys. Lett.* **B422** (1998) 247; *Nucl. Phys,.* **B537** (1999) 443.

[7]. T. Schäfer and F. Wilczek, *Phys. Rev. Lett.* **82** (1999) 3956; *Phys. Lett.* **B450** (1999) 325.

[8]. R. Rapp, T. Schäfer, E. Shuryak and M. Velkovsky, *Phys. Rev. Lett.*, **81** (1998) 53.

[9]. N.Evans, S. Hsu and M. Schwetz, *Nucl. Phys.* **B551** (1999) 275; *Phys. Lett.* **B449** (1999) 281.

[10]. D. T. Son, *Phys. Rev.* **D59** (1998) 094019.

[11]. G. W. Carter and D. Diakonov, *Phys. Rev.* **D60** (1999) 01004.

[12]. R. Rapp, T. Schäfer, E. Shuryak and M. Velkovsky, preprint IASSNS-HEP-99/40, Princeton, 1999, hep-ph/9904353.

[13]. A. Chodos, H. Minakata and F. Cooper, *Phys. Lett.* **B449** (1999) 260.

[14]. R. D. Pisarski and D. H. Rischke, *Phys. Rev.* **D60** (1999) 094013.

[15]. T Schäfer and F. Wilczek, *Phys. Rev.* **D60** (1999) 074014, 114033.

[16]. V. A. Miransky, I. A. Shovkovy and L. C. Wijewardhana, *Phys. Lett.* **B468** (1999) 270.

[17]. R. Casalbuoni and R. Gatto, *Phys. Lett.* **B464** (1999) 111.

[18]. D. K. Hong, M. Rho and I. Zahed, *Phys. Lett.* **B468** (1999) 261.

[19]. M. Rho, E. Shuryak, A. Wirzba and I. Zahed, *hep-/0001104.*

[20]. C. Manuel and M. H. G. Tytgat, *hep-ph/0001095.*

[21]. W. Brown, J. T. Liu and H. C. Ren, *hep-ph/0003199.*

[22]. J. Berges and K. Rajagopal, *Nucl. Phys.* **B358** (1999) 215.

[23]. M. Harada and A. Shibata, *Phys. Rev.* **D59** (1998) 014010.

[24]. R. D. Pisarski and D. H. Rischke, *Phys. Rev. Lett.* **83** (1999) 37.

[25]. Nguyen Van Hieu, *Basics of Fuctional Integral Technique in Quantum Field Theory of Many-Body Systems, VNUH Pub.*, Hanoi, 1999.

[26]. Nguyen Van Hieu and Le Trong Tuong, *Commun. Phys.* **8** (1998) 129.

[27]. Nguyen Van Hieu, Nguyen Hung Son, Ngo Van Thanh and Hoang Ba Thang, *Adv . Nat. Sci.* **1** (2000) 61, *hep-ph/0001251.*

[28]. Nguyen van Hieu, Hoang Ngoc Long, Nguyen Hung Son and Nguyen Ai Viet, *Proc. 24th Conference in Theoretical Physics, Samson 19-21 August 1999, p.1*, *hep-ph/0001234.*

[29]. C. Wetterich, *hep-ph/0008150.*